

( . 2).

)

- 0,1\_0,3.

- 3-4 .

10%,

- 2,

(IUCN SSC Invasive Species Specialist Group, ISSG).

(Global Invasive

Species Database, GISD).

GISD

33

(21

), GISD

(9 )

(3 ).

23

Facility, GBIF). IUCN (Global Biodiversity Information Facility, GBIF) 130  
 1272 GISD. GBIF  
 (https://www.gbif.org/ru/dataset/b351a324-77c4-41c9-a909-f30f77268bc4).

GBIF, (http://www.sevin.ru/top100worst/priortargets.html).

sevin.ru 13 : *Anadara kagoshimensis* (Tokunaga, 1906), *Arcuatula senhousia* (Benson, 1842), *Corbicula fluminea* (O.F. Muller, 1774), *Crassostrea gigas* (Thunberg, 1793), *Dreissena bugensis* (Andrusov, 1897), *Dreissena polymorpha* (Pallas, 1771), *Lithoglyphus naticoides* (C. Pfeiffer, 1828), *Magallana gigas* (Thunberg, 1793), *Mytilopsis leucophaeata* (Conrad, 1831), *Potamopyrgus antipodarum* (J.E. Gray, 1843), *Rangia cuneata* (J.E. Gray, 1843), *Rapana venosa* (Valenciennes, 1846), *Teredo navalis* Linnaeus, 1758.

GenBank.

**(PYRRHOCORIS APTERUS L.)**

• • , • • , • • , • • , [ , 2007].