

To the theory of anisotropic plane elasticity

Alexandre Soldatov

Dedicated to Professor Heinrich Begehr on the occasion of his 70th birthday

Summary: The Lamé system of general anisotropic plane elasticity is considered. A representation of a general solution of the system through a so-called Douglis analytic functions is given. The cases of orthotropic and isotropic media are also considered.

1 Lamé system

The stress tensor

$$\sigma = \begin{pmatrix} \sigma_1 & \sigma_3 \\ \sigma_3 & \sigma_2 \end{pmatrix}$$

of plane elasticity medium is connected with a displacement vector $u = \downarrow (u_1, u_2)$ by the Hook law [1]

$$\begin{pmatrix} \sigma_1 \\ \sigma_3 \end{pmatrix} = a_{11} \frac{\partial u}{\partial x} + a_{12} \frac{\partial u}{\partial y}, \quad \begin{pmatrix} \sigma_3 \\ \sigma_2 \end{pmatrix} = a_{21} \frac{\partial u}{\partial x} + a_{22} \frac{\partial u}{\partial y}. \quad (1.1)$$

The coefficients $a_{ij} \in \mathbb{R}^{2 \times 2}$ are defined by

$$\begin{aligned} a_{11} &= \begin{pmatrix} \alpha_1 & \alpha_6 \\ \alpha_6 & \alpha_3 \end{pmatrix}, & a_{12} &= \begin{pmatrix} \alpha_6 & \alpha_4 \\ \alpha_3 & \alpha_5 \end{pmatrix}, \\ a_{21} &= \begin{pmatrix} \alpha_6 & \alpha_3 \\ \alpha_4 & \alpha_5 \end{pmatrix}, & a_{22} &= \begin{pmatrix} \alpha_3 & \alpha_5 \\ \alpha_5 & \alpha_2 \end{pmatrix}, \end{aligned} \quad (1.2)$$

where modulus elasticity α_j form the positively defined matrix

$$\alpha = \begin{pmatrix} \alpha_1 & \alpha_4 & \alpha_6 \\ \alpha_4 & \alpha_2 & \alpha_5 \\ \alpha_6 & \alpha_5 & \alpha_3 \end{pmatrix}.$$

AMS 1991 subject classification: Primary: 35J55; Secondary: 30E25

Key words and phrases: Elliptic systems, anisotropic plane elasticity, Douglis analytic functions, explicit representations

By the Silvester criterium we have

$$\alpha_j > 0, \quad j = 1, 2, 3, \quad \alpha_1\alpha_2 > \alpha_4^2, \quad (1.3a)$$

$$\alpha_1\alpha_2\alpha_3 + 2\alpha_4\alpha_5\alpha_6 > \alpha_1\alpha_5^2 + \alpha_2\alpha_6^2 + \alpha_3\alpha_4^2. \quad (1.3b)$$

The elastic medium is orthotropic if $\alpha_5 = \alpha_6 = 0$, and is isotropic if

$$\alpha_5 = \alpha_6 = 0, \quad \alpha_1 = \alpha_2 = 2\alpha_3 + \alpha_4. \quad (1.4)$$

The stress tensor satisfies the equilibrium equation

$$\frac{\partial}{\partial x} \begin{pmatrix} \sigma_1 \\ \sigma_3 \end{pmatrix} + \frac{\partial}{\partial y} \begin{pmatrix} \sigma_3 \\ \sigma_2 \end{pmatrix} = 0.$$

Together with (1.1) it yields the Lamé system

$$a_{11} \frac{\partial^2 u}{\partial x^2} + (a_{12} + a_{21}) \frac{\partial^2 u}{\partial y^2} + a_{22} \frac{\partial^2 u}{\partial y^2} = 0 \quad (1.5)$$

for the displacement vector $u = (u_1, u_2)$. Besides there exists a so-called conjugate function $v(x, y)$ determined by the following relations:

$$\begin{pmatrix} \sigma_1 \\ \sigma_3 \end{pmatrix} = -\frac{\partial v}{\partial y}, \quad \begin{pmatrix} \sigma_3 \\ \sigma_2 \end{pmatrix} = \frac{\partial v}{\partial x}. \quad (1.6)$$

According to (1.1) this function is connected with u by relations

$$\frac{\partial v}{\partial x} = -\left(a_{21} \frac{\partial u}{\partial x} + a_{22} \frac{\partial u}{\partial y}\right), \quad \frac{\partial v}{\partial y} = a_{11} \frac{\partial u}{\partial x} + a_{12} \frac{\partial u}{\partial y}. \quad (1.7)$$

From (1.2) it follows that the matrix

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \in \mathbb{R}^{4 \times 4}$$

is symmetric and nonnegatively defined i.e. $(A\xi, \xi) \geq 0$ for all $\xi \in \mathbb{R}^4$. Moreover

$$(A\xi, \xi) = 0 \Leftrightarrow A\xi = 0 \Leftrightarrow \xi = (0, t, -t, 0), t \in \mathbb{R}.$$

Hence

$$(p(t)\xi_0, \xi_0) = (a_{11}\xi_0 + a_{12}t\xi_0, \xi_0) + (a_{21}\xi_0 + a_{22}t\xi_0, t\xi_0) > 0$$

for all $t \in \mathbb{R}$ and $\xi_0 \in \mathbb{R}^2$, where $p(z) = a_{11} + (a_{12} + a_{21})z + a_{22}z^2$. In particular the Lamé system is strongly elliptic [2] and its characteristic equation $\chi(z) = \det p(z)$ has no real roots. Thus for the set σ_+ of these roots in the upper half-plane we have only the following two possibilities

$$(i) \sigma_+ = \{\nu_1, \nu_2\}, \nu_1 \neq \nu_2, \quad (ii) \sigma_+ = \{\nu\}. \quad (1.8)$$

In the explicit form we have

$$p = \begin{pmatrix} p_1 & p_3 \\ p_3 & p_2 \end{pmatrix}, \quad \begin{aligned} p_1(z) &= \alpha_1 + 2\alpha_6 z + \alpha_3 z^2, \\ p_2(z) &= \alpha_3 + 2\alpha_5 z + \alpha_2 z^2, \\ p_3(z) &= \alpha_6 + (\alpha_3 + \alpha_4)z + \alpha_5 z^2, \end{aligned} \quad \chi = p_1 p_2 - p_3^2. \quad (1.9)$$

The roots of the characteristic equation can be calculated explicitly in the orthotropic case. In this case

$$\chi(z) = \alpha_2 \alpha_3 (\rho^4 + 2m\rho^2 z^2 + z^4), \quad \rho = \sqrt[4]{\frac{\alpha_1}{\alpha_2}}, \quad m = \frac{\alpha_1 \alpha_2 - \alpha_4^2 - 2\alpha_3 \alpha_4}{2\alpha_3 \sqrt{\alpha_1 \alpha_2}}.$$

It is obvious

$$m + 1 = \frac{m_1}{2\alpha_3 \sqrt{\alpha_1 \alpha_2}}, \quad m - 1 = \frac{m_2}{2\alpha_3 \sqrt{\alpha_1 \alpha_2}} (\sqrt{\alpha_1 \alpha_2} - 2\alpha_3 - \alpha_4),$$

where $m_1 = (\sqrt{\alpha_1 \alpha_2} - \alpha_4)(\sqrt{\alpha_1 \alpha_2} + \alpha_4 + 2\alpha_3)$ and $m_2 = \sqrt{\alpha_1 \alpha_2} + \alpha_4$. By virtue of (1.3a) these numbers are positive. From this equation it follows that

$$\nu_1 = i\rho e^{i\theta}, \quad \nu_2 = i\rho e^{-i\theta}, \quad \sqrt{\alpha_1 \alpha_2} < 2\alpha_3 + \alpha_4, \quad 2\theta = \arccos r, \quad (1.10a)$$

$$\nu_1 = i\rho e^t, \quad \nu_2 = i\rho e^{-t}, \quad \sqrt{\alpha_1 \alpha_2} > 2\alpha_3 + \alpha_4, \quad 2t = \operatorname{arcch} r, \quad (1.10b)$$

$$\nu_1 = \nu_2 = i\rho, \quad \sqrt{\alpha_1 \alpha_2} = 2\alpha_3 + \alpha_4, \quad (1.10c)$$

Very simple expressions we have in the case

$$\alpha_5 = \alpha_6 = 0, \quad \alpha_3 + \alpha_4 = 0. \quad (1.11)$$

Then Lamé system is diagonal and

$$\nu_1 = i\sqrt{\frac{\alpha_1}{\alpha_3}}, \quad \nu_2 = i\sqrt{\frac{\alpha_3}{\alpha_2}}. \quad (1.12)$$

This corresponds to (1.10b) with

$$m = \frac{\alpha_1 \alpha_2 + \alpha_3^2}{2\alpha_3 \sqrt{\alpha_1 \alpha_2}}.$$

The second possibility (ii) of multiple roots is corresponds to (1.10c). The equality $\rho = 1$ is valid if and only if the orthotropic medium is isotropic.

For general anisotropic Lamé system let us consider a case when three elements of the matrix $p(\nu)$ are equal to zero.

Lemma 1.1 (a) *The equalities $p_2(\nu) = p_3(\nu) = 0$, $\nu \in \sigma_+$, hold if and only if*

$$\alpha_3^2 < \alpha_1 \alpha_1, \quad |\alpha_5| < \alpha_2, \quad \alpha_3 \alpha_5 = \alpha_2 \alpha_6, \quad \alpha_2(\alpha_3 + \alpha_4) = 2\alpha_5^2. \quad (1.13a)$$

(b) *The equalities $p_1(\nu) = p_3(\nu) = 0$, $\nu \in \sigma_+$, hold if and only if*

$$\alpha_3^2 < \alpha_1 \alpha_1, \quad |\alpha_6| < \alpha_3, \quad \alpha_1 \alpha_5 = \alpha_3 \alpha_6, \quad \alpha_1(\alpha_3 + \alpha_4) = 2\alpha_6^2. \quad (1.13b)$$

(c) The both conditions (1.13) are equivalent to (1.11).

(d) The equalities $p_1(\nu) = p_2(\nu) = p_3(\nu) = 0$ are impossible for all ν . The equalities $p_2(\nu) = p_3(\nu) = 0$ or $p_1(\nu) = p_3(\nu) = 0$ are only possible in the case (i).

Proof:

(a) The equalities $p_2(\nu) = p_3(\nu) = 0$ are equivalent to the relation $p_3 = \lambda p_2$ for some $\lambda \in \mathbb{R}$, i.e.

$$\alpha_6 = \lambda \alpha_3, \quad \alpha_5 = \lambda \alpha_2 \quad \alpha_3 + \alpha_4 = 2\lambda \alpha_5 = 2\lambda^2 \alpha_2. \quad (1.14)$$

By virtue of (1.3)

$$\alpha_3 - \sqrt{\alpha_1 \alpha_2} < 2\lambda^2 \alpha_2 < \alpha_3 + \sqrt{\alpha_1 \alpha_2} \quad (1.15)$$

and

$$\alpha_1 \alpha_2 \alpha_3 + 2(2\lambda^2 \alpha_2 - \alpha_3) \lambda^2 \alpha_2 \alpha_3 > (\alpha_1 \alpha_2^2 + \alpha_2 \alpha_3^2) \lambda^2 + \alpha_3 (2\lambda^2 \alpha_2 - \alpha_3)^2.$$

The last inequality can be written in the form $(\lambda^2 \alpha_2 - \alpha_3)(\alpha_3^2 - \alpha_1 \alpha_2) > 0$. The inequalities $\lambda^2 \alpha_2 - \alpha_3 > 0$ and $\alpha_3^2 - \alpha_1 \alpha_2 > 0$ contradict to (1.15), so $\lambda^2 \alpha_2 - \alpha_3 < 0$ and $\alpha_3^2 - \alpha_1 \alpha_2 < 0$. In this case (1.15) hold automatically and we receive (1.13a) after illuminating the parameter λ from (1.14).

(b) The proof is analogously to (a).

(c) Suppose that (1.13a) and (1.13b) hold but $\alpha_3 + \alpha_4 \neq 0$. Then $\alpha_5 \alpha_6 \neq 0$ and from the system $\alpha_3 \alpha_5 - \alpha_2 \alpha_6 = 0$, $\alpha_1 \alpha_5 - \alpha_3 \alpha_6 = 0$ it follows that $\alpha_1 \alpha_2 = \alpha_3^2$. But this equality contradicts to (1.13).

(d) The first assertion follows from (c). Suppose further that for example $p_2(\nu) = p_3(\nu) = 0$ for the multiple root ν . Then $\chi(\nu) = \chi'(\nu) = 0$. As $\chi' = p_1' p_2 + p_1 p_2' - 2p_3 p_3'$ and $p_i'(\nu) \neq 0$, $i = 1, 2$, for all ν , $\text{Im } \nu \neq 0$. So we receive $p_2(\nu) = 0$ and $p_1(\nu) = p_2(\nu) = p_3(\nu) = 0$, that is impossible. \square

2 Function theoretic approach

The classic approach to plane elasticity is based [3] on representation of general solution of the Lamé system through two analytic functions. In the isotropic case this representation is known as Kolosov–Muskhelishvili formula [4]. Later there were developed various function theoretic methods [5, 6, 7], where the role of analytic functions play solutions of first order elliptic systems. Our approach to plane elasticity is based [8, 9] on the so called Douglis analytic functions which satisfy by definition the following system

$$\frac{\partial \phi}{\partial y} - J \frac{\partial \phi}{\partial x} = 0.$$

At this point the spectrum $\sigma(J)$ of the matrix $J \in \mathbb{C}^{2 \times 2}$ here coincides with σ_+ and there exists the matrix $b \in \mathbb{C}^{2 \times 2}$ such that

$$a_{11}b + (a_{12} + a_{21})bJ + a_{22}bJ^2 = 0, \quad \det \begin{pmatrix} b & \bar{b} \\ bJ & \bar{b}J \end{pmatrix} \neq 0. \quad (2.1)$$

In this terms a general solution u of the Lamé system and its conjugate function v can be represented by formulas

$$u = \operatorname{Re} b\phi, \quad v = \operatorname{Re} c\phi + \xi,$$

where $\xi \in \mathbb{R}^2$ and $c = -(a_{21}b + a_{22}bJ)$.

The matrix b can be chosen in a Jordan form. In the case (ii) by virtue of Lemma 1.1(d) the matrix J doesn't have to be equal to scalar matrix ν . So according to (1.8) there are two possibilities

$$(i) J = \begin{pmatrix} \nu_1 & 0 \\ 0 & \nu_2 \end{pmatrix}, \quad (ii) J = \begin{pmatrix} \nu & 1 \\ 0 & \nu \end{pmatrix}. \quad (2.2)$$

The matrix b is not uniquely defined by (2.1). If \tilde{b} satisfies the same conditions and $\tilde{c} = -(a_{21}\tilde{b} + a_{22}\tilde{b}J)$, then we have [10]

$$\tilde{b} = bd, \quad \tilde{c} = cd, \quad (2.3)$$

where an invertible matrix d according two cases (i) and (ii) has a form

$$(i) d = \begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix}, \quad (ii) d = \begin{pmatrix} d_1 & d_2 \\ 0 & d_1 \end{pmatrix}.$$

The matrixes b and c were described in [8, 9]. In this paper we give another more exact expressions for these matrixes. Let us introduce the matrixes

$$q = \begin{pmatrix} p_2 & -p_3 \\ -p_3 & p_1 \end{pmatrix}, \quad r(z) = -(a_{21} + a_{22}z)q(z). \quad (2.4)$$

In the explicit form

$$r(z) = \begin{pmatrix} -zq_3 & -q_1 \\ q_3 & q_2 - zq_3 \end{pmatrix}, \quad \begin{aligned} q_1(z) &= \beta_2 - \beta_5z + \beta_4z^2, \\ q_2(z) &= \beta_5 - \beta_3z + \beta_6z^2, \\ q_3(z) &= \beta_4 - \beta_6z + \beta_1z^2, \end{aligned} \quad (2.5)$$

where

$$\begin{aligned} \beta_1 &= \alpha_2\alpha_3 - \alpha_5^2, & \beta_2 &= \alpha_1\alpha_3 - \alpha_6^2, & \beta_3 &= \alpha_1\alpha_2 - \alpha_4^2, \\ \beta_4 &= \alpha_5\alpha_6 - \alpha_3\alpha_4, & \beta_5 &= \alpha_4\alpha_6 - \alpha_1\alpha_5, & \beta_6 &= \alpha_4\alpha_5 - \alpha_2\alpha_6. \end{aligned}$$

Note that β_j coincide with elements of 3×3 -matrix β , which is adjoint to the matrix α , i.e.

$$\beta = (\det \alpha)\alpha^{-1} = \begin{pmatrix} \beta_1 & \beta_4 & \beta_6 \\ \beta_4 & \beta_2 & \beta_5 \\ \beta_6 & \beta_5 & \beta_3 \end{pmatrix}. \quad (2.6)$$

Theorem 2.1 (i) Let $\sigma_+ = \{\nu_1, \nu_2\}$. If the condition (1.13a) doesn't valid then

$$b = \begin{pmatrix} p_2(\nu_1) & p_2(\nu_2) \\ -p_3(\nu_1) & -p_3(\nu_2) \end{pmatrix}, \quad c = \begin{pmatrix} -\nu_1 q_3(\nu_1) & -\nu_2 q_3(\nu_2) \\ q_3(\nu_1) & q_3(\nu_2) \end{pmatrix}. \quad (2.7a)$$

If these conditions hold and $\alpha_3 + \alpha_4 \neq 0$, then

$$b = \begin{pmatrix} -p_3(\nu_1) & p_3(\nu_2) \\ p_1(\nu_1) & p_1(\nu_2) \end{pmatrix}, \quad c = \begin{pmatrix} -q_1(\nu_1) & -q_1(\nu_2) \\ q_2(\nu_1) - \nu_1 q_3(\nu_1) & q_2(\nu_2) - \nu_2 q_3(\nu_2) \end{pmatrix}. \quad (2.7b)$$

At last in the case (1.11) we can put

$$b = 1, \quad c = -(a_{21} + a_{22}J) = \begin{pmatrix} -\alpha_3 \nu_1 & -\alpha_3 \\ \alpha_3 & -\alpha_2 \nu_2 \end{pmatrix}. \quad (2.7c)$$

(ii) Let $\sigma_+ = \{\nu\}$. Then we can put

$$b = \begin{pmatrix} p_2(\nu) & p_2'(\nu) \\ -p_3(\nu) & -p_3'(\nu) \end{pmatrix}, \quad c = \begin{pmatrix} -\nu q_3(\nu) & -q_3(\nu) - \nu q_3'(\nu) \\ q_3(\nu) & q_3'(\nu) \end{pmatrix}. \quad (2.8)$$

Proof:

- (i) From (2.1) it follows that the columns $b_{(k)}$, $k = 1, 2$, satisfy the equation $p(\nu_k)b_{(k)} = 0$. Taking into account (2.4) we have $p(z)q(z) = \chi(z)$ and hence $p(\nu_k)q_{(i)}(\nu_k) = 0$, $i = 1, 2$. So we can put $b_{(k)} = d_k q_{(i)}(\nu_k)$, $d_k \neq 0$, under assumption $q_{(i)}(\nu_k) \neq 0$. If the conditions (1.13a) have no place then then according to Lemma 1.1 this assumption is fulfilled for $i = 1$.

Let the conditions (1.13a) hold. Then the unit matrix $b = 1$ satisfies (2.1) in the case (1.11). If $\alpha_3 + \alpha_4 \neq 0$, then by lemma 1 we have $b_{(k)} = d_k q_{(2)}(\nu_k)$, $d_k \neq 0$ for all $k = 1, 2$. By virtue of (2.3) we can put here $d_1 = d_2 = 1$.

Let turn to the matrix $c = -(a_{21}b + a_{22}bJ)$. It is obviously that $c_{(k)} = -a_{21}b_{(k)} - \nu_k a_{22}b_{(k)}$ and therefore

$$c_{(k)} = -(a_{21} + a_{22}\nu_k)p_{(i)}(\nu_k), \quad b_{(k)} = p_{(i)}(\nu_k).$$

Taking into account (2.3) we complete the proof.

- (ii) It follows from (2.1) that

$$p(\nu)b_{(1)} = 0, \quad p(\nu)b_{(2)} + p'(\nu)b_{(1)} = 0.$$

Since the root ν is multiple we have $p(\nu)q'(\nu)p'(\nu)q(\nu) = 0$. By virtue of Lemma 1.1 the column $q_{(1)}(\nu) \neq 0$ and therefore we can write

$$b_{(1)} = d_1 q_{(1)}(\nu), \quad b_{(2)} = d_1 q_{(1)}'(\nu) + d_2 q_{(1)}(\nu)$$

with $d_1 \neq 0$. Taking into account (2.3) we complete the proof for the matrix b .

As $(bJ)_1 = \nu b_1$, $(bJ)_2 = b_1 + \nu b_2$, we can write

$$c_{(1)} = -a_{21}b_{(1)} - \nu a_{22}b_{(1)}, \quad c_{(2)} = -a_{21}b_{(2)} - \nu a_{22}b_{(2)} - a_{22}b_{(1)}.$$

Putting $b_{(1)} = p_{(1)}(\nu)$, $b_{(2)} = p'_{(1)}(\nu)$ we receive

$$c_{(1)} = r_{(1)}(\nu), \quad c_{(2)} = -(a_{21} + \nu a_{22})p'_{(1)}(\nu) - a_{22}p_{(1)}(\nu) = r'_{(1)}(\nu),$$

that complete the proof. \square

Due to [10] the matrix b is invertible for all strong elliptic system and in particular for Lamé system. The matrix c has the same property.

Theorem 2.2 *Under assumptions of the Theorem 2.1 the matrix c is invertible.*

Proof: Within notations (2.6) the characteristic polynomial $\chi = p_1p_2 - p_3^2$ can be written in the form

$$\chi(z) = q_1(z) - zq_2(z) + z^2q_3(z). \quad (2.9)$$

The expressions (2.5) for q_j yield the relation $\xi = \beta\eta$ with respect to the vectors $\xi = (q_3, q_1, q_2)$ and $\eta = (z^2, 1, -z)$. Taking into account (2.6) we conclude that $(\det \alpha)\eta = \alpha\xi$ or

$$\begin{aligned} (\det \alpha)z^2 &= \alpha_4q_1 + \alpha_6q_2 + \alpha_1q_3, \\ \det \alpha &= \alpha_2q_1 + \alpha_5q_2 + \alpha_4q_3, \\ -(\det \alpha)z &= \alpha_5q_1 + \alpha_3q_2 + \alpha_6q_3. \end{aligned} \quad (2.10)$$

In particular the common equalities $q_1(\nu) = q_2(\nu) = q_3(\nu) = 0$ are impossible for all ν . From this and (2.9) it follows that only one of numbers $q_i(\nu)$, $i = 1, 2, 3$, where $\nu \in \sigma_+$, may be equal to zero.

The following implications

$$q_3(\nu) = 0 \quad \Leftrightarrow \quad p_2(\nu) = p_3(\nu) = 0, \quad (2.11a)$$

$$q_1(\nu) = 0 \quad \Rightarrow \quad p_3(\nu) = 0. \quad (2.11b)$$

for every $\nu \in \sigma_+$ hold.

In fact let $\chi(\nu) = q_3(\nu) = 0$. Then by virtue of (2.9) we can write $q_1(\nu) = \lambda\nu$, $q_2(\nu) = \lambda \neq 0$, $q_3(\nu) = 0$. Putting $z = \nu$ in (2.10) we conclude that $\alpha_3 + 2\alpha_5\nu + \alpha_2\nu^2 = 0$. Accordingly (1.9) this expression coincides with $p_2(\nu) = 0$. Since $p_1(\nu)p_2(\nu) - p_3^2(\nu) = 0$ we have also $p_3(\nu) = 0$. Conversely if $p_2(\nu) = p_3(\nu) = 0$, then by virtue of (2.4), (2.5) $q_3(\nu) = 0$.

The second implication (2.11b) is proved analogously. If $\chi(\nu) = q_1(\nu) = 0$, then $q_1 = 0$, $q_2 = \lambda\nu$, $q_3 = \lambda$ and we derive from (2.10) that $p_3(\nu) = 0$.

Let the conditions (1.13a) be broken. Then by virtue of (2.11a) we have $q_2(\nu) \neq 0$ for $\nu \in \sigma_+$ and it is easily verified that $\det c \neq 0$ in the cases (2.7a) and (2.8). Let the conditions (1.13a) hold and therefor $p_2(\nu) = p_3(\nu) = 0$ for some $\nu \in \sigma_+$. For definiteness let $\nu = \nu_1$. Then by virtue of (2.11a) $q_3(\nu_1) = 0$ and therefor $q_1(\nu_1) \neq 0$.

Let us prove that also $q_1(\nu_2) \neq 0$ out of the exceptional case (1.11). Really if $q_1(\nu_1) = 0$ then according to (2.11b) we will have $p_3(\nu_2) = 0$. As $p_2(\nu_2) \neq 0$ it follows from the equality $p_1(\nu)p_2(\nu) - p_3^2(\nu) = 0$ that $p_1(\nu_2) = 0$. So $p_2(\nu) = p_3(\nu) = 0$, $\nu = \nu_1$ and $p_1(\nu) = p_3(\nu) = 0$, $\nu = \nu_2$ and by virtue of Lemma 1.1(a), (b) the both conditions (1.13) hold which is equivalent to (1.11).

Thus the numbers $q_1(\nu_j)$ in (2.7b) are not equal to zero. It follows from (2.9) that $q_2(\nu_j) - \nu_j q_3(\nu_j) = \nu_j^{-1} q_1(\nu_j)$ and so $\det c \neq 0$.

According to (2.2c) in the exceptional case (1.11) we have $\det c = \alpha_3^2 + \alpha_2 \alpha_3 \nu_1 \nu_2$. Taking into account (1.12) we also receive $\det c \neq 0$.

The expressions of the Theorem 2.1 are simplified in the orthotropic case. In this case (1.9) and (2.5) have the form

$$\begin{aligned} p_1(z) &= \alpha_1 + \alpha_3 z^2, & p_2(z) &= \alpha_3 + \alpha_2 z^2, \\ p_3(z) &= (\alpha_3 + \alpha_4)z, & q_3(z) &= -\alpha_3(\alpha_4 - \alpha_2 z^2). \end{aligned}$$

If $\alpha_3 + \alpha_4 \neq 0$ then we can use the formulas (2.7a) and (2.8). So we have the expressions

$$b = \begin{pmatrix} \alpha_3 + \alpha_2 \nu_1^2 & \alpha_3 + \alpha_2 \nu_2^2 \\ -(\alpha_3 + \alpha_4)\nu_1 & -(\alpha_3 + \alpha_4)\nu_2 \end{pmatrix}, \quad c = \alpha_3 \begin{pmatrix} \nu_1(\alpha_4 - \alpha_2 \nu_1^2) & \nu_2(\alpha_4 - \alpha_2 \nu_2^2) \\ -(\alpha_4 - \alpha_2 \nu_1^2) & -(\alpha_4 - \alpha_2 \nu_2^2) \end{pmatrix},$$

where ν_j are defined by (1.10a) or (1.10b), and

$$b = \begin{pmatrix} \alpha_3 - \alpha_2 \rho^2 & 2i\alpha_2 \rho \\ -i(\alpha_3 + \alpha_4)\rho & -(\alpha_3 + \alpha_4) \end{pmatrix}, \quad c = \alpha_3 \begin{pmatrix} i\rho(\alpha_4 + \alpha_2 \rho^2) & \alpha_4 + 3\alpha_2 \rho^2 \\ -(\alpha_4 + \alpha_2 \rho^2) & 2i\alpha_2 \rho \end{pmatrix}$$

in the case (ii).

The last formulas permit further simplification in the isotropic case. In this case $\rho = 1$ and $\alpha_1 > \alpha_3$ by virtue of (1.3a), (1.4). So

$$b = \begin{pmatrix} \alpha_3 - \alpha_1 & 2\alpha_1 i \\ (\alpha_3 - \alpha_1)i & \alpha_3 - \alpha_1 \end{pmatrix}, \quad c = 2\alpha_3 \begin{pmatrix} (\alpha_1 - \alpha_3)i & 2\alpha_1 - \alpha_3 \\ \alpha_3 - \alpha_1 & \alpha_1 i \end{pmatrix}.$$

According to (2.3) we can multiply these matrices by

$$d = (\alpha_3 - \alpha_1)^{-1} \begin{pmatrix} 1 & 2\alpha_1(\alpha_1 - \alpha_3)^{-1}i \\ 0 & 1 \end{pmatrix}.$$

As a result we have

$$b = \begin{pmatrix} 1 & 0 \\ i & -\mathfrak{x} \end{pmatrix}, \quad c = \alpha_3 \begin{pmatrix} 2i & \mathfrak{x} - 1 \\ 2 & i(\mathfrak{x} + 1) \end{pmatrix},$$

where $\mathfrak{x} = (\alpha_1 + \alpha_3)/(\alpha_1 - \alpha_3)$. □

3 Conjugate function

Let us consider a second order elliptic system

$$a_{11} \frac{\partial^2 u}{\partial x^2} + a_{(12)} \frac{\partial^2 u}{\partial y^2} + a_{22} \frac{\partial^2 u}{\partial y^2} = 0, \quad a_{(12)} = a_{12} + a_{21}, \quad (3.1)$$

with coefficients $a_{ij} \in \mathbb{R}^{l \times l}$. We can introduce the notion of conjugate function v to solution $u = (u_1, \dots, u_l)$ of this equation as above by (1.7). Of course this definition depends on the partition $a_{(12)} = a_{12} + a_{21}$. There is a question which second order system defines the function v ? Let us put

$$a_1 = a_{11}^{-1}a_{12}, \quad a_2 = a_{22}^{-1}a_{21} \quad (3.2)$$

and define matrixes $d_1, d_2 \in \mathbb{R}^{l \times l}$ such that

$$d_1 a_{22} (1 - a_2 a_1) = d_2 a_{11} (1 - a_1 a_2). \quad (3.3)$$

Lemma 3.1 *The conjugate function v satisfies the system*

$$d_1 \frac{\partial^2 v}{\partial x^2} + (d_1 a_{21} a_{11}^{-1} + d_2 a_{12} a_{22}^{-1}) \frac{\partial^2 v}{\partial x \partial y} + d_2 \frac{\partial^2 v}{\partial y^2} = 0. \quad (3.4)$$

Proof: With respect to the vectors

$$U = \left(\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y} \right), \quad V = \left(\frac{\partial v}{\partial x}, \frac{\partial v}{\partial y} \right)$$

(1.7) takes a form

$$V = \begin{pmatrix} -a_{21} & -a_{22} \\ a_{11} & a_{12} \end{pmatrix} U. \quad (3.5)$$

Hence (3.1) and the analogous equation

$$d_{11} \frac{\partial^2 u}{\partial x^2} + d_{(12)} \frac{\partial^2 u}{\partial y^2} + d_{22} \frac{\partial^2 u}{\partial y^2} = 0 \quad (3.6)$$

for the function v can be rewritten as

$$\frac{\partial U}{\partial y} = \begin{pmatrix} 0 & 1 \\ -a_{22}^{-1} a_{11} & -a_{22}^{-1} a_{(12)} \end{pmatrix} \frac{\partial U}{\partial x}, \quad \begin{pmatrix} 1 & 0 \\ 0 & d_{22} \end{pmatrix} \frac{\partial V}{\partial y} = \begin{pmatrix} 0 & 1 \\ -d_{11} & -d_{(12)} \end{pmatrix} \frac{\partial V}{\partial x}.$$

Together with (3.5) it follows that

$$\left[\begin{pmatrix} 1 & 0 \\ 0 & d_{22} \end{pmatrix} \begin{pmatrix} -a_{21} & -a_{22} \\ a_{11} & a_{12} \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -a_{22}^{-1} a_{11} & -a_{22}^{-1} a_{(12)} \end{pmatrix} - \begin{pmatrix} 0 & 1 \\ -d_{11} & -d_{(12)} \end{pmatrix} \begin{pmatrix} -a_{21} & -a_{22} \\ a_{11} & a_{12} \end{pmatrix} \right] \frac{\partial U}{\partial x} = 0$$

for all U . This is equivalent to system

$$d_{11} a_{21} - d_{(12)} a_{11} = -d_{22} (a_{12} a_{22}^{-1} a_{11}), \quad d_{11} a_{22} - d_{(12)} a_{12} = d_{22} (a_{11} - a_{12} a_{22}^{-1} a_{(12)})$$

with respect to unknown coefficients $d_{11}, d_{22} \in \mathbb{R}^{l \times l}$. This system we can rewrite as following:

$$d_{22} (a_{11} - a_{12} a_{22}^{-1} a_{21}) = d_{11} (a_{22} - a_{21} a_{11}^{-1} a_{12}), \quad d_{(12)} = d_{11} a_{21} a_{11}^{-1} + d_{22} a_{12} a_{22}^{-1}.$$

The first equation coincides with (3.3) with respect to $d_i = d_{ii}$, but a substitution of second one to (3.6) gives (3.4).

Let us apply this result to Lamé system. According to (2) the matrices (3.2) have the form

$$a_1 = \frac{1}{\alpha_1\alpha_3 - \alpha_6^2} \begin{pmatrix} 0 & \alpha_3\alpha_4 - \alpha_5\alpha_6 \\ \alpha_1\alpha_3 - \alpha_6^2 & \alpha_1\alpha_5 - \alpha_4\alpha_6 \end{pmatrix} = \begin{pmatrix} 0 & -\beta_4/\beta_2 \\ 1 & -\beta_5/\beta_2 \end{pmatrix},$$

$$a_2 = \frac{1}{\alpha_2\alpha_3 - \alpha_5^2} \begin{pmatrix} \alpha_2\alpha_6 - \alpha_4\alpha_5 & \alpha_2\alpha_3 - \alpha_5^2 \\ \alpha_3\alpha_4 - \alpha_5\alpha_6 & 0 \end{pmatrix} = \begin{pmatrix} -\beta_6/\beta_1 & 1 \\ -\beta_4/\beta_1 & 0 \end{pmatrix},$$

where β_j figure in (2.5), (2.6). Analogously

$$a_{21}a_{11}^{-1} = \begin{pmatrix} 0 & 1 \\ -\beta_4/\beta_2 & -\beta_5/\beta_2 \end{pmatrix}, \quad a_{12}a_{22}^{-1} = \begin{pmatrix} -\beta_6/\beta_1 & -\beta_4/\beta_1 \\ 1 & 0 \end{pmatrix}.$$

Hence

$$1 - a_1a_2 = \frac{1}{\beta_1\beta_2} \begin{pmatrix} \beta_1\beta_2 - \beta_4^2 & 0 \\ \beta_2\beta_6 - \beta_4\beta_5 & 0 \end{pmatrix}, \quad 1 - a_2a_1 = \frac{1}{\beta_1\beta_2} \begin{pmatrix} 0 & \beta_1\beta_5 - \beta_4\beta_6 \\ 0 & \beta_1\beta_2 - \beta_4^2 \end{pmatrix}$$

and (3.3) reduces to conditions $(d_1)_{(2)} = (d_2)_{(1)}$ with respect to columns of the matrixes d_j . So we can take

$$d_1 = \begin{pmatrix} s_1 & 0 \\ t_1 & 0 \end{pmatrix}, \quad d_2 = \begin{pmatrix} 0 & s_2 \\ 0 & t_2 \end{pmatrix}$$

with some $s_j, t_j \in \mathbb{R}$ and (3.4) reduces to

$$\begin{pmatrix} s_1 & 0 \\ t_1 & 0 \end{pmatrix} \frac{\partial^2 v}{\partial x^2} + \begin{pmatrix} s_2 & s_1 \\ t_2 & t_1 \end{pmatrix} \frac{\partial^2 v}{\partial x \partial y} + \begin{pmatrix} 0 & s_2 \\ 0 & t_2 \end{pmatrix} \frac{\partial^2 v}{\partial y^2} = 0.$$

This system is equivalent to equations

$$\frac{\partial}{\partial x} \left(\frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} \right) = 0, \quad \frac{\partial}{\partial y} \left(\frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} \right) = 0.$$

But they are consequence of the equation (1.6) from which it follows

$$\frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} = 0.$$

Therefore the result of Lemma 3.1 for the Lamé system is reduced to the last equation. \square

Acknowledgements. The work was supported in part by the Russian Foundation of Basic Research (RFBR)(project No. 07-01-00299) and by the National Natural Science Foundation of China (NSFC) in the framework of the bilateral project "Complex Analysis and its applications" (project No. 08-01-92208-GFEN)

References

- [1] V. D. Kupradze. *Potential Methods in the Theory of Elasticity*, Israel Program Sc. Transl., Jerusalem, 1965.
- [2] M. I. Vishik, *On strong elliptic systems of differential equations*, *Matematicheskii Sbornik*, 29: 615–676, 1951, (in Russian).
- [3] S. G. Lekhnitskii. *Anisotropic Plates*, Gordon and Breach, New York, 1968.
- [4] N. I. Muskhelishvili, *Some Basic Problems of the Mathematical Theory of Elasticity*, Noordhoff, Groningen, 1953.
- [5] H. Begehr, Lin Wei, *A mixed–contact problem in orthotropic elasticity*, in *Partial differential equations with Real Analysis*, H. Begerhand, A. Jeffrey, eds., Longman Scientific & Technical, 1992, 219–239.
- [6] R. P. Gilbert. *Plane Ellipticity and Related Problems*, Amer. Math. Soc., Providence, RI, 1981.
- [7] R. P. Gilbert, Lin Wei. *Function theoretical solutions to problems of orthotropic elasticity*, *J. Elasticity*, 15: 143–154, 1985.
- [8] A. P. Soldatov. *Lame system of anisotropic plane elasticity*, *Dokl. RAN*, 385: 163–167, 2002 (in Russian).
- [9] A. P. Soldatov. *On elliptic boundary value problems on upper half-plane*, *J. of Applied Functional Analysis*, 1: 85–98, 2007.
- [10] A. P. Soldatov *On the first and second boundary value problems for elliptic systems on the plane*, *Differential'nye Uravneniya*, 39: 674–686, 2003 (in Russian).

Alexandre Soldatov
Belgorod State University
Pobeda 85
Belgorod, 308015
Russia
soldatov@bsu.edu.ru
soldatov48@mail.ru